arXiv:1203.2212v1 [math.CA] 9 Mar 2012

A symmetric Nörlund sum with application to inequalities

Artur M. C. Brito da Cruz, Natália Martins and Delfim F. M. Torres

[Submitted 18-Oct-2011; accepted 09-March-2012; to Proceedings of International Conference on Differential & Difference Equations and Applications, in honour of Professor Ravi P. Agarwal, to be published by Springer in the series *Proceedings in Mathematics* (PROM)]

Abstract Properties of an α,β-symmetric Nörlund sum are studied. Inspired in the work by Agarwal et al., α,β-symmetric quantum versions of Hölder, Cauchy–Schwarz and Minkowski inequalities are obtained.

1 Introduction

The symmetric derivative of function f at point x is defined as $\lim_{h\to 0}(f(x+h)-f(x-h))/(2h)$. The notion of symmetrically differentiable is interesting because if a function is differentiable at a point then it is also symmetrically differentiable, but the converse is not true. The best known example of this fact is the absolute value function: $f(x)=|x|$ is not differentiable at $x=0$ but is symmetrically differentiable at $x=0$ with symmetric derivative zero [6].

Quantum calculus is, roughly speaking, the equivalent to traditional infinitesimal calculus but without limits [4]. Therefore, one can introduce the symmetric quantum derivative of f at x by $(f(x+h)-f(x-h))/(2h)$. As in any calculus, it is then natural to develop a corresponding integration theory, looking to such integral as the inverse operator of the derivative.

Artur M. C. Brito da Cruz[1,2]
e-mail: `artur.cruz@estsetubal.ips.pt`

Natália Martins[2]
e-mail: `natalia@ua.pt`

Delfim F. M. Torres[2]
e-mail: `delfim@ua.pt`

[1]Escola Superior de Tecnologia de Setúbal, Estefanilha, 2910-761 Setúbal, Portugal

[2]Center for Research and Development in Mathematics and Applications
Department of Mathematics, University of Aveiro, 3810-193 Aveiro, Portugal

The main goal of this paper is to study the properties of a general symmetric quantum integral that we call, due to the so-called Nörlund sum [4], the α,β-symmetric Nörlund sum.

The paper is organized as follows. In Section 2 we define the forward and backward Nörlund sums. Then, in Section 3, we introduce the α,β-symmetric Nörlund sum and give some of its properties. We end with Section 4, proving α,β-symmetric versions of Hölder's, Cauchy–Schwarz's and Minkowski's inequalities.

2 Forward and backward Nörlund sums

This section is dedicated to the inverse operators of the α-forward and β-backward differences, $\alpha > 0$, $\beta > 0$, defined respectively by

$$\Delta_\alpha [f](t) := \frac{f(t+\alpha) - f(t)}{\alpha}, \quad \nabla_\beta [f](t) := \frac{f(t) - f(t-\beta)}{\beta}.$$

Definition 1. Let $I \subseteq \mathbb{R}$ be such that $a, b \in I$ with $a < b$ and $\sup I = +\infty$. For $f : I \to \mathbb{R}$ and $\alpha > 0$ we define the Nörlund sum (the α-forward integral) of f from a to b by

$$\int_a^b f(t)\,\Delta_\alpha t = \int_a^{+\infty} f(t)\,\Delta_\alpha t - \int_b^{+\infty} f(t)\,\Delta_\alpha t,$$

where $\int_x^{+\infty} f(t)\,\Delta_\alpha t = \alpha \sum_{k=0}^{+\infty} f(x+k\alpha)$, provided the series converges at $x = a$ and $x = b$. In that case, f is said to be α-forward integrable on $[a,b]$. We say that f is α-forward integrable over I if it is α-forward integrable for all $a, b \in I$.

Until Definition 2 (the backward/nabla case), we assume that I is an interval of $\mathbb{R}$ such that $\sup I = +\infty$. Note that if $f : I \to \mathbb{R}$ is a function such that $\sup I < +\infty$, then we can extend function f to $\tilde{f} : \tilde{I} \to \mathbb{R}$, where $\tilde{I}$ is an interval with $\sup \tilde{I} = +\infty$, in the following way: $\tilde{f}|_I = f$ and $\tilde{f}|_{\tilde{I} \setminus I} = 0$.

Using the techniques of Aldwoah in his Ph.D. thesis [2], it can be proved that the α-forward integral has the following properties:

Theorem 1. *If $f, g : I \to \mathbb{R}$ are α-forward integrable on $[a,b]$, $c \in [a,b]$, $k \in \mathbb{R}$, then*

1. $\int_a^a f(t)\,\Delta_\alpha t = 0$;
2. $\int_a^b f(t)\,\Delta_\alpha t = \int_a^c f(t)\,\Delta_\alpha t + \int_c^b f(t)\,\Delta_\alpha t$, *when the integrals exist;*
3. $\int_a^b f(t)\,\Delta_\alpha t = -\int_b^a f(t)\,\Delta_\alpha t$;
4. kf *is α-forward integrable on $[a,b]$ and* $\int_a^b kf(t)\,\Delta_\alpha t = k\int_a^b f(t)\,\Delta_\alpha t$;
5. $f+g$ *is α-forward integrable on $[a,b]$ and*

$$\int_a^b (f+g)(t)\,\Delta_\alpha t = \int_a^b f(t)\,\Delta_\alpha t + \int_a^b g(t)\,\Delta_\alpha t;$$

6. *if* $f \equiv 0$, *then* $\int_a^b f(t)\,\Delta_\alpha t = 0$.

Theorem 2. *Let* $f : I \to \mathbb{R}$ *be* α*-forward integrable on* $[a,b]$. *If* $g : I \to \mathbb{R}$ *is a nonnegative* α*-forward integrable function on* $[a,b]$, *then* fg *is* α*-forward integrable on* $[a,b]$.

Proof. Since g is α-forward integrable, then both series $\alpha\sum_{k=0}^{+\infty} g(a+k\alpha)$ and $\alpha\sum_{k=0}^{+\infty} g(b+k\alpha)$ converge. We want to study the nature of series $\alpha\sum_{k=0}^{+\infty} fg(a+k\alpha)$ and $\alpha\sum_{k=0}^{+\infty} fg(b+k\alpha)$. Since there exists an order $N \in \mathbb{N}$ such that $|fg(b+k\alpha)| \leqslant g(b+k\alpha)$ and $|fg(a+k\alpha)| \leqslant g(a+k\alpha)$ for all $k > N$, then both $\alpha\sum_{k=0}^{+\infty} fg(a+k\alpha)$ and $\alpha\sum_{k=0}^{+\infty} fg(b+k\alpha)$ converge absolutely. The intended conclusion follows.

Theorem 3. *Let* $f : I \to \mathbb{R}$ *and* $p > 1$. *If* $|f|$ *is* α*-forward integrable on* $[a,b]$, *then* $|f|^p$ *is also* α*-forward integrable on* $[a,b]$.

Proof. There exists $N \in \mathbb{N}$ such that $|f(b+k\alpha)|^p \leqslant |f(b+k\alpha)|$ and $|f(a+k\alpha)|^p \leqslant |f(a+k\alpha)|$ for all $k > N$. Therefore, $|f|^p$ is α-forward integrable on $[a,b]$.

Theorem 4. *Let* $f,g : I \to \mathbb{R}$ *be* α*-forward integrable on* $[a,b]$. *If* $|f(t)| \leqslant g(t)$ *for all* $t \in \{a+k\alpha : k \in \mathbb{N}_0\}$, *then for* $b \in \{a+k\alpha : k \in \mathbb{N}_0\}$ *one has*

$$\left|\int_a^b f(t)\,\Delta_\alpha t\right| \leqslant \int_a^b g(t)\,\Delta_\alpha t.$$

Proof. Since $b \in \{a+k\alpha : k \in \mathbb{N}_0\}$, there exists k_1 such that $b = a+k_1\alpha$. Thus,

$$\begin{aligned}\left|\int_a^b f(t)\,\Delta_\alpha t\right| &= \left|\alpha\sum_{k=0}^{+\infty} f(a+k\alpha) - \alpha\sum_{k=0}^{+\infty} f(a+(k_1+k)\alpha)\right| \\ &= \left|\alpha\sum_{k=0}^{+\infty} f(a+k\alpha) - \alpha\sum_{k=k_1}^{+\infty} f(a+k\alpha)\right| = \left|\alpha\sum_{k=0}^{k_1-1} f(a+k\alpha)\right| \\ &\leqslant \alpha\sum_{k=0}^{k_1-1} |f(a+k\alpha)| \leqslant \alpha\sum_{k=0}^{k_1-1} g(a+k\alpha) \\ &= \alpha\sum_{k=0}^{+\infty} g(a+k\alpha) - \alpha\sum_{k=k_1}^{+\infty} g(a+k\alpha) = \int_a^b g(t)\,\Delta_\alpha t.\end{aligned}$$

Corollary 1. *Let* $f,g : I \to \mathbb{R}$ *be* α*-forward integrable on* $[a,b]$ *with* $b = a+k\alpha$ *for some* $k \in \mathbb{N}_0$.

1. *If* $f(t) \geqslant 0$ *for all* $t \in \{a+k\alpha : k \in \mathbb{N}_0\}$, *then* $\int_a^b f(t)\,\Delta_\alpha t \geqslant 0$.
2. *If* $g(t) \geqslant f(t)$ *for all* $t \in \{a+k\alpha : k \in \mathbb{N}_0\}$, *then* $\int_a^b g(t)\,\Delta_\alpha t \geqslant \int_a^b f(t)\,\Delta_\alpha t$.

We can now prove the following fundamental theorem of the α-forward calculus.

Theorem 5 (Fundamental theorem of Nörlund calculus). *Let* $f : I \to \mathbb{R}$ *be* α-*forward integrable over* I. *Let* $x \in I$ *and define* $F(x) := \int_a^x f(t)\,\Delta_\alpha t$. *Then,* $\Delta_\alpha[F](x) = f(x)$. *Conversely,* $\int_a^b \Delta_\alpha[f](t)\,\Delta_\alpha t = f(b) - f(a)$.

Proof. If $G(x) = -\int_x^{+\infty} f(t)\,\Delta_\alpha t$, then

$$\begin{aligned}\Delta_\alpha[G](x) &= \frac{G(x+\alpha)-G(x)}{\alpha} = \frac{-\alpha\sum_{k=0}^{+\infty} f(x+\alpha+k\alpha)+\alpha\sum_{k=0}^{+\infty} f(x+k\alpha)}{\alpha} \\ &= \sum_{k=0}^{+\infty} f(x+k\alpha) - \sum_{k=0}^{+\infty} f(x+(k+1)\alpha) = f(x).\end{aligned}$$

Therefore, $\Delta_\alpha[F](x) = \Delta_\alpha\left(\int_a^{+\infty} f(t)\,\Delta_\alpha t - \int_x^{+\infty} f(t)\,\Delta_\alpha t\right) = f(x)$. Using the definition of α-forward difference operator, the second part of the theorem is also a consequence of the properties of Mengoli's series. Since

$$\begin{aligned}\int_a^{+\infty} \Delta_\alpha[f](t)\,\Delta_\alpha t &= \alpha\sum_{k=0}^{+\infty} \Delta_\alpha[f](a+k\alpha) = \alpha\sum_{k=0}^{+\infty} \frac{f(a+k\alpha+\alpha)-f(a+k\alpha)}{\alpha} \\ &= \sum_{k=0}^{+\infty}\Big(f(a+(k+1)\alpha) - f(a+k\alpha)\Big) = -f(a)\end{aligned}$$

and $\int_b^{+\infty} \Delta_\alpha[f](t)\,\Delta_\alpha t = -f(b)$, it follows that

$$\int_a^b \Delta_\alpha[f](t)\,\Delta_\alpha t = \int_a^{+\infty} f(t)\,\Delta_\alpha t - \int_b^{+\infty} f(t)\,\Delta_\alpha t = f(b) - f(a).$$

Corollary 2 (α-forward integration by parts). *Let* $f, g : I \to \mathbb{R}$. *If* fg *and* $f\Delta_\alpha[g]$ *are* α-*forward integrable on* $[a,b]$, *then*

$$\int_a^b f(t)\,\Delta_\alpha[g](t)\,\Delta_\alpha t = f(t)\,g(t)\Big|_a^b - \int_a^b \Delta_\alpha[f](t)\,g(t+\alpha)\,\Delta_\alpha t$$

Proof. Since $\Delta_\alpha[fg](t) = \Delta_\alpha[f](t)\,g(t+\alpha) + f(t)\,\Delta_\alpha[g](t)$, then

$$\begin{aligned}\int_a^b f(t)\,\Delta_\alpha[g](t)\,\Delta_\alpha t &= \int_a^b \Big(\Delta_\alpha[fg](t) - \Delta_\alpha[f](t)\,g(t+\alpha)\Big)\Delta_\alpha t \\ &= \int_a^b \Delta_\alpha[fg](t)\,\Delta_\alpha t - \int_a^b \Delta_\alpha[f](t)\,g(t+\alpha)\,\Delta_\alpha t \\ &= f(t)\,g(t)\Big|_a^b - \int_a^b \Delta_\alpha[f](t)\,g(t+\alpha)\,\Delta_\alpha t.\end{aligned}$$

Remark 1. Our study of the Nörlund sum is in agreement with the Hahn quantum calculus [2,3,5]. In [4] $\int_a^b f(t)\,\Delta_\alpha t = \alpha\,[f(a) + f(a+\alpha) + \cdots + f(b-\alpha)]$ for $a < b$ such that $b - a \in \alpha\mathbb{Z}$, $\alpha \in \mathbb{R}^+$. In contrast with [4], our definition is valid for any two real points a, b and not only for those points belonging to the time scale $\alpha\mathbb{Z}$. The definitions (only) coincide if function f is α-forward integrable on $[a,b]$.

Similarly, we introduce the β-backward integral.

Definition 2. Let I be an interval of $\mathbb{R}$ such that $a,b \in I$ with $a<b$ and $\inf I = -\infty$. For $f : I \to \mathbb{R}$ and $\beta > 0$ we define the β-backward integral of f from a to b by

$$\int_a^b f(t)\nabla_\beta t = \int_{-\infty}^b f(t)\nabla_\beta t - \int_{-\infty}^a f(t)\nabla_\beta t,$$

where $\int_{-\infty}^x f(t)\nabla_\beta t = \beta \sum_{k=0}^{+\infty} f(x-k\beta)$, provided the series converges at $x=a$ and $x=b$. In that case, f is called β-backward integrable on $[a,b]$. We say that f is β-backward integrable over I if it is β-backward integrable for all $a,b \in I$.

The β-backward Nörlund sum has similar results and properties as the α-forward Nörlund sum. In particular, the β-backward integral is the inverse operator of ∇_β.

3 The α,β-symmetric Nörlund sum

We define the α,β-symmetric integral as a linear combination of the α-forward and the β-backward integrals.

Definition 3. Let $f : \mathbb{R} \to \mathbb{R}$ and $a,b \in \mathbb{R}$, $a<b$. If f is α-forward and β-backward integrable on $[a,b]$, $\alpha,\beta \geq 0$ with $\alpha+\beta>0$, then we define the α,β-symmetric integral of f from a to b by

$$\int_a^b f(t)\, d_{\alpha,\beta}t = \frac{\alpha}{\alpha+\beta}\int_a^b f(t)\Delta_\alpha t + \frac{\beta}{\alpha+\beta}\int_a^b f(t)\nabla_\beta t.$$

Function f is α,β-symmetric integrable if it is α,β-symmetric integrable for all $a,b \in \mathbb{R}$.

Remark 2. Note that if $\alpha \in \mathbb{R}^+$ and $\beta = 0$, then $\int_a^b f(t)\, d_{\alpha,\beta}t = \int_a^b f(t)\Delta_\alpha t$ and we do not need to assume in Definition 3 that f is β-backward integrable; if $\alpha = 0$ and $\beta \in \mathbb{R}^+$, then $\int_a^b f(t)\, d_{\alpha,\beta}t = \int_a^b f(t)\nabla_\beta t$ and we do not need to assume that f is α-forward integrable.

Example 1. Let $f(t) = 1/t^2$. Then $\int_1^3 \frac{1}{t^2} d_{2,2}t = \frac{10}{9}$.

The α,β-symmetric integral has the following properties:

Theorem 6. *Let $f,g : \mathbb{R} \to \mathbb{R}$ be α,β-symmetric integrable on $[a,b]$. Let $c \in [a,b]$ and $k \in \mathbb{R}$. Then,*

1. $\int_a^a f(t)\, d_{\alpha,\beta}t = 0$;

2. $\int_a^b f(t)\,d_{\alpha,\beta}t = \int_a^c f(t)\,d_{\alpha,\beta}t + \int_c^b f(t)\,d_{\alpha,\beta}t$*, when the integrals exist;*

3. $\int_a^b f(t)\,d_{\alpha,\beta}t = -\int_b^a f(t)\,d_{\alpha,\beta}t$*;*

4. kf *is* α,β*-symmetric integrable on* $[a,b]$ *and* $\int_a^b kf(t)\,d_{\alpha,\beta}t = k\int_a^b f(t)\,d_{\alpha,\beta}t$*;*

5. $f+g$ *is* α,β*-symmetric integrable on* $[a,b]$ *and*

$$\int_a^b (f+g)(t)\,d_{\alpha,\beta}t = \int_a^b f(t)\,d_{\alpha,\beta}t + \int_a^b g(t)\,d_{\alpha,\beta}t;$$

6. fg *is* α,β*-symmetric integrable on* $[a,b]$ *provided* g *is a nonnegative function.*

Proof. These results are easy consequences of the α-forward and β-backward integral properties.

The next result follows immediately from Theorem 3 and the corresponding β-backward version.

Theorem 7. *Let* $f:\mathbb{R}\to\mathbb{R}$ *and* $p>1$. *If* $|f|$ *is symmetric* α,β*-integrable on* $[a,b]$, *then* $|f|^p$ *is also* α,β*-symmetric integrable on* $[a,b]$.

Theorem 8. *Let* $f,g:\mathbb{R}\to\mathbb{R}$ *be* α,β*-symmetric integrable functions on* $[a,b]$, $\mathscr{A} := \{a+k\alpha : k\in\mathbb{N}_0\}$ *and* $\mathscr{B} := \{b-k\beta : k\in\mathbb{N}_0\}$. *For* $b\in\mathscr{A}$ *and* $a\in\mathscr{B}$ *one has:*

1. if $|f(t)| \leqslant g(t)$ *for all* $t\in\mathscr{A}\cup\mathscr{B}$, *then* $\left|\int_a^b f(t)\,d_{\alpha,\beta}t\right| \leqslant \int_a^b g(t)\,d_{\alpha,\beta}t$*;*

2. if $f(t)\geqslant 0$ *for all* $t\in\mathscr{A}\cup\mathscr{B}$, *then* $\int_a^b f(t)\,d_{\alpha,\beta}t \geqslant 0$*;*

3. if $g(t)\geqslant f(t)$ *for all* $t\in\mathscr{A}\cup\mathscr{B}$, *then* $\int_a^b g(t)\,d_{\alpha,\beta}t \geqslant \int_a^b f(t)\,d_{\alpha,\beta}t$.

Proof. It follows from Theorem 4 and Corollary 1 and the corresponding β-backward versions.

In Theorem 9 we assume that $a,b\in\mathbb{R}$ with $b\in\mathscr{A} := \{a+k\alpha : k\in\mathbb{N}_0\}$ and $a\in\mathscr{B} := \{b-k\beta : k\in\mathbb{N}_0\}$, where $\alpha,\beta\in\mathbb{R}_0^+$, $\alpha+\beta\neq 0$.

Theorem 9 (Mean value theorem). *Let* $f,g:\mathbb{R}\to\mathbb{R}$ *be bounded and* α,β*-symmetric integrable on* $[a,b]$ *with* g *nonnegative. Let* m *and* M *be the infimum and the supremum, respectively, of function* f. *Then, there exists a real number* K *satisfying the inequalities* $m\leqslant K\leqslant M$ *such that* $\int_a^b f(t)g(t)\,d_{\alpha,\beta}t = K\int_a^b g(t)\,d_{\alpha,\beta}t$.

Proof. Since $m\leqslant f(t)\leqslant M$ for all $t\in\mathbb{R}$ and $g(t)\geqslant 0$, then $mg(t)\leqslant f(t)g(t)\leqslant Mg(t)$ for all $t\in\mathscr{A}\cup\mathscr{B}$. All functions mg, fg and Mg are α,β-symmetric integrable on $[a,b]$. By Theorems 6 and 8, $m\int_a^b g(t)\,d_{\alpha,\beta}t \leqslant \int_a^b f(t)g(t)\,d_{\alpha,\beta}t \leqslant M\int_a^b g(t)\,d_{\alpha,\beta}t$. If $\int_a^b g(t)\,d_{\alpha,\beta}t = 0$, then $\int_a^b f(t)g(t)\,d_{\alpha,\beta}t = 0$; if $\int_a^b g(t)\,d_{\alpha,\beta}t > 0$, then $m\leqslant \frac{\int_a^b f(t)g(t)d_{\alpha,\beta}t}{\int_a^b g(t)d_{\alpha,\beta}t}\leqslant M$. Therefore, the middle term of these inequalities is equal to a number K, which yields the intended result.

4 α,β-Symmetric Integral Inequalities

Inspired in the work by Agarwal et al. [1], we now present α,β-symmetric versions of Hölder, Cauchy–Schwarz and Minkowski inequalities. As before, we assume that $a,b \in \mathbb{R}$ with $b \in \mathscr{A} := \{a + k\alpha : k \in \mathbb{N}_0\}$ and $a \in \mathscr{B} := \{b - k\beta : k \in \mathbb{N}_0\}$, where $\alpha,\beta \in \mathbb{R}_0^+$, $\alpha + \beta \neq 0$.

Theorem 10 (Hölder's inequality). *Let $f,g : \mathbb{R} \to \mathbb{R}$ and $a,b \in \mathbb{R}$ with $a < b$. If $|f|$ and $|g|$ are α,β-symmetric integrable on $[a,b]$, then*

$$\int_a^b |f(t)g(t)|\, d_{\alpha,\beta}t \leqslant \left(\int_a^b |f(t)|^p\, d_{\alpha,\beta}t\right)^{\frac{1}{p}} \left(\int_a^b |g(t)|^q\, d_{\alpha,\beta}t\right)^{\frac{1}{q}}, \tag{1}$$

where $p > 1$ and $q = p/(p-1)$.

Proof. For $\alpha,\beta \in \mathbb{R}_0^+$, $\alpha + \beta \neq 0$, the following inequality holds: $\alpha^{\frac{1}{p}}\beta^{\frac{1}{q}} \leqslant \frac{\alpha}{p} + \frac{\beta}{q}$. Without loss of generality, suppose that $\left(\int_a^b |f(t)|^p\, d_{\alpha,\beta}t\right)\left(\int_a^b |g(t)|^q\, d_{\alpha,\beta}t\right) \neq 0$ (note that both integrals exist by Theorem 7). Set $\xi(t) = |f(t)|^p / \int_a^b |f(\tau)|^p\, d_{\alpha,\beta}\tau$ and $\gamma(t) = |g(t)|^q / \int_a^b |g(\tau)|^q\, d_{\alpha,\beta}\tau$. Since both functions α and β are symmetric α,β-integrable on $[a,b]$, then (1) holds:

$$\begin{aligned}
\int_a^b \frac{|f(t)|}{\left(\int_a^b |f(\tau)|^p\, d_{\alpha,\beta}\tau\right)^{\frac{1}{p}}} \frac{|g(t)|}{\left(\int_a^b |g(\tau)|^q\, d_{\alpha,\beta}\tau\right)^{\frac{1}{q}}} d_{\alpha,\beta}t &= \int_a^b \xi(t)^{\frac{1}{p}}\gamma(t)^{\frac{1}{q}}\, d_{\alpha,\beta}t \\
&\leqslant \int_a^b \left(\frac{\xi(t)}{p} + \frac{\gamma(t)}{q}\right) d_{\alpha,\beta}t \\
&= \frac{1}{p}\int_a^b \left(\frac{|f(t)|^p}{\int_a^b |f(\tau)|^p\, d_{\alpha,\beta}\tau}\right) d_{\alpha,\beta}t + \frac{1}{q}\int_a^b \left(\frac{|g(t)|^q}{\int_a^b |g(\tau)|^q\, d_{\alpha,\beta}\tau}\right) d_{\alpha,\beta}t = 1.
\end{aligned}$$

The particular case $p = q = 2$ of (1) gives the Cauchy–Schwarz inequality.

Corollary 3 (Cauchy–Schwarz's inequality). *Let $f,g : \mathbb{R} \to \mathbb{R}$ and $a,b \in \mathbb{R}$ with $a < b$. If f and g are α,β-symmetric integrable on $[a,b]$, then*

$$\int_a^b |f(t)g(t)|\, d_{\alpha,\beta}t \leqslant \sqrt{\left(\int_a^b |f(t)|^2\, d_{\alpha,\beta}t\right)\left(\int_a^b |g(t)|^2\, d_{\alpha,\beta}t\right)}.$$

We prove the Minkowski inequality using Hölder's inequality.

Theorem 11 (Minkowski's inequality). *Let $f,g : \mathbb{R} \to \mathbb{R}$ and $a,b,p \in \mathbb{R}$ with $a < b$ and $p > 1$. If f and g are α,β-symmetric integrable on $[a,b]$, then*

$$\left(\int_a^b |f(t)+g(t)|^p\, d_{\alpha,\beta}t\right)^{\frac{1}{p}} \leqslant \left(\int_a^b |f(t)|^p\, d_{\alpha,\beta}t\right)^{\frac{1}{p}} + \left(\int_a^b |g(t)|^p\, d_{\alpha,\beta}t\right)^{\frac{1}{p}}.$$

Proof. One has

$$\begin{aligned}\int_a^b |f(t)+g(t)|^p d_{\alpha,\beta}t &= \int_a^b |f(t)+g(t)|^{p-1}|f(t)+g(t)|\, d_{\alpha,\beta}t \\ &\leqslant \int_a^b |f(t)|\,|f(t)+g(t)|^{p-1} d_{\alpha,\beta}t + \int_a^b |g(t)|\,|f(t)+g(t)|^{p-1} d_{\alpha,\beta}t.\end{aligned}$$

Applying Hölder's inequality (Theorem 10) with $q = p/(p-1)$, we obtain

$$\begin{aligned}&\int_a^b |f(t)+g(t)|^p d_{\alpha,\beta}t \leqslant \left(\int_a^b |f(t)|^p d_{\alpha,\beta}t\right)^{\frac{1}{p}} \left(\int_a^b |f(t)+g(t)|^{(p-1)q} d_{\alpha,\beta}t\right)^{\frac{1}{q}} \\ &+ \left(\int_a^b |g(t)|^p d_{\alpha,\beta}t\right)^{\frac{1}{p}} \left(\int_a^b |f(t)+g(t)|^{(p-1)q} d_{\alpha,\beta}t\right)^{\frac{1}{q}} \\ &= \left[\left(\int_a^b |f(t)|^p d_{\alpha,\beta}t\right)^{\frac{1}{p}} + \left(\int_a^b |g(t)|^p d_{\alpha,\beta}t\right)^{\frac{1}{p}}\right] \left(\int_a^b |f(t)+g(t)|^{(p-1)q} d_{\alpha,\beta}t\right)^{\frac{1}{q}}.\end{aligned}$$

Therefore,

$$\frac{\int_a^b |f(t)+g(t)|^p d_{\alpha,\beta}t}{\left(\int_a^b |f(t)+g(t)|^{(p-1)q} d_{\alpha,\beta}t\right)^{\frac{1}{q}}} \leqslant \left(\int_a^b |f(t)|^p d_{\alpha,\beta}t\right)^{\frac{1}{p}} + \left(\int_a^b |g(t)|^p d_{\alpha,\beta}t\right)^{\frac{1}{p}}.$$

Our α,β-symmetric calculus is more general than the standard h-calculus. In particular, all our results give, as corollaries, results in the classical quantum h-calculus by choosing $\alpha = h > 0$ and $\beta = 0$.

Acknowledgements Work supported by FEDER and Portuguese funds, COMPETE reference FCOMP-01-0124-FEDER-022690, and CIDMA and FCT, project PEst-C/MAT/UI4106/2011. Brito da Cruz is also supported by FCT through the Ph.D. fellowship SFRH/BD/33634/2009.

References

1. R. Agarwal, M. Bohner and A. Peterson, Inequalities on time scales: a survey, Math. Inequal. Appl. **4** (2001), no. 4, 535–557.
2. K. A. Aldwoah, *Generalized time scales and associated difference equations*, Ph.D. thesis, Cairo University, 2009.
3. A. M. C. Brito da Cruz, N. Martins and D. F. M. Torres, Higher-order Hahn's quantum variational calculus, Nonlinear Anal. **75** (2012), no. 3, 1147–1157. `arXiv:1101.3653`
4. V. Kac and P. Cheung, *Quantum calculus*, Springer, New York, 2002.
5. A. B. Malinowska and D. F. M. Torres, The Hahn quantum variational calculus, J. Optim. Theory Appl. **147** (2010), no. 3, 419–442. `arXiv:1006.3765`
6. B. S. Thomson, *Symmetric properties of real functions*, Dekker, New York, 1994.